

## Rearing Record Chart for Hy-Line Brown (Alternative Systems)

Rearer:
House Number:


Hatch day/date:     /     /
Delivery day/date:     /     /
Total chicks delivered:

Flock Code	No. of Birds	Flock Code	No. of Birds

Age Wks	Daily Mortality and Culls							Mortality			Light Hours	Temp		Feed Delivery Tonnes		Feed Intake			Bodyweight		
								Week	Cum	Cum %		Min/Max	Week	Cum	GMS Day	Cumulative		Target (g)	Actual (g)	Evenness %	
																Actual (g)	Target (g) week				
1															14-15		98 - 105	70 - 75			
2																17-21		217 - 252	110 - 125		
3																23-25		378 - 427	170 - 190		
4																28-30		574 - 637	240 - 270		
5																34-36		812 - 889	330 - 360		
6																38-40		1078 - 1169	420 - 465		
7																42-44		1372 - 1477	520 - 560		
8																53-57		1743 - 1876	620 - 670		
9																55-59		2128 - 2289	730 - 780		
10																58-62		2534 - 2723	840 - 890		
11																62-65		2968 - 3178	940 - 990		
12																65-69		3423 - 3661	1030 - 1080		
13																69-72		3906 - 4165	1110 - 1165		
14																74-78		4424 - 4711	1190 - 1230		
15																76-80		4956 - 5271	1260 - 1300		
16																78-82		5502 - 5845	1330 - 1370		
17																79-83		6055 - 6426	1400 - 1440		
18																					

PLEASE KEEP ACCURATE RECORDS  
An important aid to management

House Number:
No. of Birds:
Day Old Date:
Date Housed:



It should be noted that at the time of transfer from rearing to laying, there will be some loss in body weight (which is normally in the region of 10-12%). This loss is mainly due to reduced water intake and some dehydration of the pullet. To help regain these losses, please refer to our management guide.

Higher target weight standards used